



EVALUATION OF NUTRIENT DIGESTIBILITY IN *LABEO ROHITA* JUVENILES FED LYSINE SUPPLEMENTED DIET

Faryad Ali ¹, Muhammad Afzal ², Syed Zakir Hussain Shah ^{*3}, Mahroze Fatima ⁴, Khadija Akram ⁵ and Rashid Hussain ⁶

^{1,5} M.Phil. Students, ² Professor, Fish Nutrition Laboratory, Department of Zoology, Wildlife and Fisheries, ⁶ Ph.D. Scholar, Institute of Horticultural Sciences, University of Agriculture, Faisalabad, ³ Assistant Professor, Department of Zoology, University of Gujrat, Gujrat, ⁴ Assistant Professor, Department of Fisheries and Aquaculture, University of Veterinary and Animal Sciences, Lahore, Pakistan.

*Corresponding author e-mail: zakiruaf@gmail.com

Article received on: 23/01/2017

Accepted for publication: 10/12/2018

ABSTRACT

A study was conducted under the laboratory conditions in the Department of Zoology, Wildlife and Fisheries, University of Agriculture, Faisalabad, Pakistan during 2014 to investigate the effect of supplementation of lysine on nutrient digestibility in the juveniles of *Labeo rohita*. Seven experimental diets were formulated by supplementing 0, 0.5, 1, 1.5, 2, 2.5 and 3% lysine and designated as D1, D2, D3, D4, D5, D6 and D7, respectively. To test each diet, two replicates with 20 fish each (average weight 3.526±0.056g) were assigned. The experiment was performed within the time duration of 60 days. Water quality parameters like dissolved oxygen (5.8-7.3 mg/L), temperature (24.9-28.7°C) and pH (7.4-8.6) were maintained throughout this period. Results showed that lysine supplementation in fish diet increased the digestibility of crude protein and dry matter. Maximum digestibility of crude protein and dry matter was observed at lysine level of 1.5 and 2.5%, respectively. However, it was observed that digestibility of crude fat decreased in response to lysine supplementation. In conclusion, it was found that lysine supplementation improved the nutrient digestibility performance of juveniles of *L. rohita*

KEYWORDS: *Labeo rohita*; Rohu; fingerlings; lysine; nutrient digestibility; Pakistan.

INTRODUCTION

The rate of utilization of fish meal in aqua feeds is increasing tremendously day by day due to increasing demand of aquaculture production (Forster and Ogata, 1998). Now-a-days, plant based meals are used to meet the protein and energy demands of aquatic animals because plant origin oils such as rice bran oil, peanut oil, soybean oil and linseed oil have good fatty acid profile which is necessary for fish (Masih *et al.*, 2014; Robinson *et al.*, 1980; Tahira and Butt, 2007). Sunflower seeds can provide Vitamin A and D, linoleic acid, oleic acid and unsaturated fatty acid (Gatlin *et al.*, 2007; Grewal and Williams, 2000). Plant based meal is the cheap source of proteins but has lower digestibility in fish meal (Carter and Hauler, 2000), To overcome this deficiency, various dietary additives have been used to improve the nutritional value of plant meals-based diets including exogenous enzymes (Shah *et al.*, 2016, 2015a, 2015b), organic acids (Akram *et al.*, 2016; Shah *et al.*, 2016; 2015c), chelated mineral (Shah and Afzal, 2016) and vitamins (Fatima Shah *et al.*, 2016).

Plant based diet is also low in available protein and has poor essential amino acid profile (Tacon, 1995). Protein being the key nutrient, is required by organisms in huge amount in diet for proper functioning of their bodies (Lovell, 1989). It is the important structural component of bones. By weight, bone constitutes 22% protein as

proteins play a key role in collagen formation. Fish require amino acid to strengthen its skeletal system (Hughes, 2003). Amino acid profile in the fish diet has an effect on dietary protein (Keembiyehetty and Gatlin, 1992). Lysine is one of the most important essential amino acid as many fish species have a considerable amount of lysine in their skeleton (Ahmed and Khan, 2004). Decreased growth and increased mortality are the major effects that can be seen in the fish fed with essential amino acid deficient diet (Ketola, 1983).

When the fish meal is replaced by the plant origin protein diet, major effect in almost all fin fish species is the deficiency of lysine in their bodies. Different fish species require different amounts of lysine in their diet ranging from 3.2 to 6.2% (Fatima *et al.*, 2016; Harris, 1980; Ozorio *et al.*, 2003; Small and Soares, 2000; Wang *et al.*, 2005; Yang *et al.*, 2010).

Moreover, lysine plays an important role in controlling the biosynthesis of carnitine which takes place in liver and skeletal muscles. Carnitine assists in the transportation of long chain unsaturated fat molecules for β -oxidation in mitochondria which can serve as a fuel during times of fasting (Tacon, 1995). Lysine also helps in the maintenance of osmotic pressure and acid base balance. It is needed for the deposition of protein in the body (Chiu *et al.*, 1987, 1988). Previous studies have shown that lysine supplementation in diet also results in the improved weight gain of fish (Hu *et al.*, 2008;

Robinson and Li, 1994; Robinson *et al.*, 1980; Viola *et al.*, 1992a; Walton *et al.*, 1984; Zarate and Lovell, 1997). The lysine requirement of fish is around 5.7% of its total dietary protein (Kim *et al.*, 1992; Anon. 1993; Singh, 1987; Wilson, 1985). Soybean meal-based diet may require supplementation of lysine due to high lysine requirement (57-70 g kg⁻¹ of protein) (Satheesha and Murthy, 1999; Ahmed and Khan, 2004).

Labeo rohita is widely cultured now-a-days due to its fast growth and disease resistance properties as it is considered as one of the tastiest fish among other fish species (Jhingran and Pullin, 1988). Accordingly, dietary lysine is essential in order to prepare nutritionally complete and cost-effective diet for intensive culture of fingerling *Labeo rohita*.

The purpose of present study was to determine the nutrient digestibility in *L. rohita* juveniles fed lysine supplemented diet.

MATERIALS AND METHODS

Labeo rohita juveniles (average weight 3.526±0.0056 g) were obtained from Government Fish Seed Hatchery, Faisalabad and transported to the Laboratory of Department of Zoology, Wildlife and Fisheries, University of Agriculture, Faisalabad, Pakistan in the year 2014. These were kept in large tanks to acclimatize with the laboratory conditions. Then these were transferred to specially designed V-shaped tanks (UA system) for the collection of fecal material from water media. Nine duplets were fed the respective test diets once in daily (Table 1). Water temperature (24.9-28.7°C), pH (7.4-8.6) and dissolved oxygen (5.8-7.3 mg/L) were monitored by the usage of thermometer, pH meter (Jenway, 3510), D.O. meter (Jenway, 970), respectively during the study period. Capillary system was used to provide aeration to all the tanks round the clock.

Table 1. Ingredients composition (%) of experimental diet.

Ingredients	Composition (%)
Fish meal	17
Sunflower meal	25
Corn gluten meal	20
Soybean meal	15
Fish oil	7
Wheat flour	5.3
Mineral mixture*	3
Vitamin mixture**	3
CaHPO ₄	2.5
Ascorbic acid	1
Chromic oxide	1
Choline chloride	0.2
Total	100

Each kg mineral granules contains Ca (Calcium) 155gm, Mn (Manganese) 2000 mg, P (Phosphorous) 135gm, Cu (Copper) 600mg, Mg (Magnesium) 55gm, Co (Cobalt) 40mg, Fe (Iron) 1000 mg, I (Iodine) 40mg, Zn (Zinc) 3000 mg, Se (Selenium) 3mg, Na (Sodium) 45gm, **Each Kg of Vitamin premix contains, Vitamin A

15 M.I.U., Vitamin D3 3 M.I.U., Nicotinic acid 25000mg, Vitamin B1 5000 mg, Vitamin E 6000 IU, Vitamin B2 6000 mg, Vitamin K3 4000 mg, Vitamin B6 4000 mg, Folic acid 750 mg, Vitamin B12 9000 mcg, Vitamin C 15000mg, Calcium pantothenate 10000mg

Experimental diets and feed ingredients

Ingredients were purchased from local poultry feed market. Prior to the formulation, chemical analysis of ingredients was carried out following AOAC (1995). Before adding in the experimental diets, the feed ingredients were ground and refined to required particle size (Table 2).

To prepare diets all dry ingredients were intermixed in electric mixer for 10-20 minutes; while mixing constantly, fish oil was gradually added during mixing. Inert marker (Chromic oxide 1%) was also incorporated during mixing. Lysine was added at the levels of 0, 0.5, 1, 1.5, 2, 2.5 and 3% to dry mixed ingredients to make seven test diets. Approximately 10-15% distilled water was added to the ingredient mixture to facilitate pelleting by lab extruder.

Table 2. Chemical composition (%) of experimental diets (dry basis).

Diet	Lysine (%)	Dry matter (%)	Crude protein (%)	Crude fat (%)
D1	0	91.905	30.849	8.7845
D2	0.5	91.565	30.9965	9.253
D3	1	92.29	31.264	9.1875
D4	1.5	92.71	30.701	9.112
D5	2	91.64	31.8945	9.7445
D6	2.5	92.225	31.194	8.9235
D7	3	91.89	31.694	8.6455

Sample collection and feeding protocol

Two replicates were assigned with stocking density of 20 fish for each test diet and diet was fed to the fish at satiation level, twice a day. Tanks were covered by net to prevent juveniles to jump out. After two hours of feeding session, the tanks were washed completely to remove particles of diet and refilled with fresh water. Fecal collection tube was used to collect faeces of each tank after two hours' gap. Fecal material of each replicated treatment dried in oven at 60°C, ground and stored for chemical analysis. The experiment lasted for 60 days.

Biochemical analysis

The samples of test diet and faeces were homogenized by pestle and mortar and analyzed by standard methods (AOAC, 1995). Oven was used to determine moisture contents by drying at 105°C for 12 hours, micro Kjeldahl apparatus for crude protein after acid digestion, petroleum ether extraction method for crude fat through Soxhlet HT2 1045 system, crude ash, by burning at 65°C or 12 hours in electric furnace (Eyela-TMF 3100) to constant weight.

Statistical analysis

The analyzed data of nutrient digestibility was subjected to statistical analysis using one way analysis of variance (ANOVA). The differences among means were tested by Newman Keuls Test and considered significant at $P < 0.05$. Costate Computer Software, Version 6.303 was used for statistical analysis.

RESULTS AND DISCUSSION

The data (Table 3) illustrate that lysine supplementation significantly improved the dry matter digestibility upto 1.5% lysine level which was decreased with further supplementation. Similarly, crude protein digestibility was also increased by dietary supplementation upto lysine level of 2.5%. However, dietary lysine supplementation showed no effect on crude fat digestibility.

Table 3. Effect of graded levels of lysine supplementation on apparent coefficient (ADC %) of nutrients in *Labeo rohita* juveniles.

Diet	Lysine level (%)	Dry matter	Crude protein	Crude fat
D1	0	72.62 ^c	89.67 ^d	85.07
D2	0.5	73.95 ^a	90.09 ^{cd}	86.32
D3	1	73.71 ^{ab}	90.39 ^c	86.36
D4	1.5	73.99 ^a	90.26 ^c	86.74
D5	2	73.26 ^b	91.19 ^b	87.98
D6	2.5	73.55 ^{ab}	91.71 ^a	87.42
D7	3	73.44 ^{ab}	90.73 ^{bc}	85.46
PSE		0.109	0.139	0.69
ANOVA				
p-value		0.0006***	0.0002***	0.1628 ns

These observations also confirm the results of El-saidy and Gaber (2002) who observed that lysine supplementations in *L. rohita* significantly ($p < 0.05$) improved the nutrient digestibility coefficients of crude protein, crude fat and gross energy with maximum increase at 0.5% L-lysine level. Similarly, Cheng *et al.* (2003) reported increased digestibility of crude

protein in *Oncorhynchus mykiss* fed with 0.2% lysine supplemented diet. According to Williams *et al.* (2001), supplementation of amino acids in low protein diets improved the growth performance of barramundi *Lates calcarifer* juveniles and that only with low protein diet, amino acids were utilized efficiently by fish. It has been observed by Zarate and Lovell (1997) that supplementation of lysine in low protein diets increased the protein and lipid contents in the edible tissue. They also found that crude fat and dry matter digestibility were not affected by lysine supplementation. Some recent studies (Li *et al.*, 2014; Tang *et al.*, 2012) showed improved nutrient digestibility and growth performance in grass carp.

In contrast to present observations, Davies *et al.* (1997) observed no significant differences in the digestibility of crude protein, crude fat and dry matter in *Oncorhynchus mykiss* fed lysine supplemented diet. Similarly, Alam *et al.* (2005) also observed non-significant effect of lysine supplementation on the nutritional attributes in *Marsupinaeus japonica*. Moreover, Biswas *et al.* (2007), did not observe any significant effect of lysine supplementation on crude protein and crude ash digestibility in *Penaeus monodon*. Researches on the channel cat fish *Ictalurus punctatus* noted that lysine or methionine had no significant effect on growth performance of the fish (Webster *et al.*, 2000).

The differences in lysine optimum levels for nutrient digestibility may be due to the differences in fish species, fish age and size, composition of basal diet, stocking density and culture conditions in the laboratory (Li *et al.*, 2014).

CONCLUSION

It is concluded that for *L. rohita* juveniles, the plant-based diet supplemented with lysine provided satisfactory nutrient digestibility performance when compared with the reference diet containing no supplementation. Hence, these results should encourage further studies on carp species for formulating low cost and more nutritive diet.




REFERENCES

- Ahmed, I. and M.A. Khan. 2004. Dietary lysine requirement of fingerling Indian major carp, *Cirrhinus mrigala* (Hamilton). *Aquaculture*. 235:499-511.
- Akram, Z., M. Afzal, S.Z.H. Shah, M. Fatima, K. Akram, S.M. Hussain and M.M. Shahzad. 2016. Role of citric acid and phytase supplementation in improving trace mineral retention in Rohu (*Labeo rohita*) juveniles fed soybean meal-based diet.

- Biologia. 62:87-93.
- Alam, M.S., S. Teshima, S. Koshio, M. Ishikawa, O. Uyan, L.H.H. Hector and R.M. Fady. 2005. Supplemental effects of coated methionine and/or lysine to soy protein isolate diet for juvenile kuruma shrimp, *Marsupenaeus japonicus*. Aquaculture. 248:13-19.
- Anon. 1993. Nutrient Requirement of Fish. National Research Council. National Academy Press, Washington DC, USA.
- AOAC, 1995. Official Methods of Analysis Association of Official Analytical Chemists, vol. 1. Cunniff, P.A. (ed.), 16th Ed. AOAC International, Arlington, USA.
- Biswas, A.K., H. Kaku, S.C. Ji, M. Seoka and K. Takii. 2007. Use of soybean meal and phytase for partial replacement of fish meal in the diet of red sea bream, *Pagrus major*. Aquaculture. 267:284-291.
- Carter, C.G. and R.C. Hauler. 2000. Fish meal replacement by plant meals in extruded feeds for Atlantic salmon, *Salmo salar*. Aquaculture. 185(3):299-311.
- Cheng, Z.J., R.W. Hardy and J.L. Usry. 2003. Effects of lysine supplementation in plant protein-based diets on the performance of rainbow trout (*Oncorhynchus mykiss*) and apparent digestibility coefficients of nutrients. Aquaculture. 215:255-265.
- Chiu, Y.N., R.E. Austic and G.L. Rumsey. 1987. Interactions among dietary minerals, arginine and lysine in rainbow trout (*Salmo gairdneri*). Fish Physiol. Biochem. 4:45-55.
- Chiu, Y.N., R.E. Austic, and G.L. Rumsey. 1988. Effect of feeding level and dietary electrolytes on the arginine requirement of rainbow trout (*Salmo gairdneri*). Aquaculture. 69(1-2):79-91.
- Davies, S.J., P.C. Morris and R.T.M. Baker. 1997. Partial substitution of fish meal and full-fat soya bean meal with wheat gluten and influence of lysine supplementation in diets for rainbow trout, *Oncorhynchus mykiss* (Walbaum). Aquacult. Res. 28(5):317-328.
- El-Saidy, D.M.S.D. and M.M.A. Gaber. 2002. Complete replacement of fish meal by soybean meal with dietary L-lysine supplementation for Nile tilapia *Oreochromis niloticus* (L.) fingerlings. J. World Aquac. Soc. 33(3):297-306.
- Fatima, M., M. Afzal, K. Abbas and M. Shahid. 2016. Effect of high vitamin E dosages on fillets lipid peroxidation of *Labeo rohita* Fingerlings. Sylwan. 160(12):191-203.
- Forster, I. and H.Y. Ogata. 1998. Lysine requirement of juvenile Japanese flounder *Paralichthys olivaceus* and juvenile red sea bream, *Pagrus major*. Aquaculture. 161:131-142.
- Gatlin, D.M., F.T. Barrows, P. Brown, K. Dabrowski, T.G. Gaylord, R.W. Hardy, E. Herman, G. Hu, A. Kroghdahl, R. Nelson, K. Overturf, M. Rust, W. Sealey, D. Skonberg, E.J. Souza, D. Stone, R. Wilson and E. Wurtele. 2007. Expanding the utilization of sustainable plant products in aqua feeds: a review. Aquacult. Res. 38(6): 551-579.
- Grewal, H.S. and R. Williams. 2000. Zinc nutrition affects alfalfa responses to water stress and excessive moisture. J. Plant Nutr. 23(7):949-962.
- Harris, L.E. 1980. Diet stuffs. In: Fish Diet Technology. Pillay, T.V.R. (ed.), UNDP/FAO, Rome, Italy. p. 111-168.
- Hu, M., Y. Wang, Q. Wang, M. Zhao, B. Xiong, X. Qian, Y. Zhao and Z. Luo. 2008. Replacement of fish meal by rendered animal protein ingredients with lysine and methionine supplementation to practical diets for gibel carp, *Carassius auratus gibelio*. Aquaculture. 275:260-265.
- Hughes, B.D. 2003. Interaction of dietary calcium and protein in bone health in humans. J. Nutr. 133(3):852-854.
- Jhingran, V.G. and R.S.V. Pullin. 1988. A hatchery manual for common, Chinese and Indian major carps under field condition. ICLARM Studies and Reviews 11. Asian Development Bank, Manila, Philippines.
- Keembiyehetty, C.N. and D.M. Gatlin. 1992. Dietary lysine requirement of juvenile hybrid striped bass (*Morone chrysops* x *M. saxatilis*). Aquaculture. 104(3-4): 271-277.
- Ketola, H.G. 1983. Requirement for dietary lysine and arginine by fry of rainbow trout. J. Anim. Sci. 56: 101-107.
- Kim, K.I., T.B. Kayesand and C.H. Amundson. 1992. Requirements for lysine and arginine by rainbow trout (*Oncorhynchus mykiss*). Aquaculture. 106(3-4):333-344.
- Li, X.Y., L. Tang, K. Hu, Y. Liu, W.D. Jiang, J. Jiang, P. Wu, G.F. Chen, S.H. Li, S.Y. Kuang and L. Feng. 2014. Effect of dietary lysine on growth, intestinal enzymes activities and antioxidant status of sub-adult grass carp (*Ctenopharyngodon idella*). Fish Physiol. Biochem. 40(3):659-671.
- Lovell, R.T. 1989. Nutrition and Feeding of Fish. Van Nostrand Reinhold, New York, NY 260 Pp. 123-234.
- Masih, S., Z. Iqbal, A.M. Arif, M. Rafiq, G. Rasool and A. Rashid. 2014. Effect of linseed oil substitution on physico-chemical properties of cookies. J. Agric. Res. 52(3):425-437.

- Ozorio, R.O.A., J.A.J. Verreth, C.R. Aragao, C.J. Vermeulen, J.W. Schrama, M.W.A. Verstegen, and E.A. Huisman. 2003. Dietary carnitine supplements increased lipid metabolism and decreased protein oxidation in African catfish (*Clarias gariepinus*) juveniles fed high fat levels. *J. Aquacult. Trop.* 18:225-238.
- Robinson, E.H. and M.H. Li. 1994. Use of plant proteins in catfish feeds: replacement of soybean meal with cottonseed meal and replacement of fish meal with soybean meal and cottonseed meal. *J. World Aquac. Soc.* 25(2):271-276.
- Robinson, E.H., R.P. Wilson and W.E. Poe. 1980. Re-evaluation of the lysine requirement and lysine utilization by fingerling channel catfish. *J. Nutr.* 110(11):2313-2316.
- Satheesha, S.K. and H.S. Murthy. 1999. Dietary requirement of Indian major carp, *Catla catla* for the essential amino acid lysine. *Indian J. Anim. Sci.* 69(4):277-279.
- Shah, S.Z.H. and M. Afzal. 2016. Effect of citric acid, phytase and organic trace elements supplementation on bone mineralization of *Labeo rohita* fingerlings fed sunflower meal-based diet. *Sylwan.* 160(12):217-231.
- Shah, S.Z.H., M. Afzal, A. Akmal, M. Fatima and S. M. Hussain. 2016. Effect of citric acid and phytase on growth performance and mineralization of *Labeo rohita* juveniles fed soybean meal-based diet. *Int. J. Agric. Biol.* 18:111-116.
- Shah, S.Z.H., M. Afzal, M. Fatima, S.M. Hussain, M.A. Bhatti and H.F. Hayat. 2015a. Supplementation of phytase and citric acid to soybean meal-based diet enhance muscle mineralization of rohu, *Labeo rohita*, juveniles. *J. Sci.* 5(9):796-800.
- Shah, S.Z.H., M. Afzal, S.M. Hussain, M. Fatima, M. Bilal, T. Ahmed and R.Z. Habib. 2015b. Supplementation of citric acid and phytase improves the digestive enzymes activities in *Labeo rohita* fingerlings. *Biologia*, 61:63-68.
- Shah, S.Z.H., M. Afzal, S.Y. Khan, S.M. Hussain and R.Z. Habib. 2015c. Prospects of using citric acid as fish feed supplement: a review. *Int. J. Agric. Biol.* 17:1-8.
- Singh, B. N. 1987. On the sulphur amino acids (methionine+cystine) and lysine requirements of rohu, *Labeo rohita* (Hamilton) and essential amino acid composition of certain promising feeds evolved for carps. *The First Indian Fisheries Forum, Mangalore, India.* p. 67.
- Small, B.C. and J.R. Soares. 2000. Quantitative dietary lysine requirement of juvenile striped bass, *Moronesaxatilis*. *Aquacult. Nutr.* 6(4):207-212.
- Tacon, A.G.J. 1995. Feed ingredients for carnivorous fish species: alternatives to fishmeal and other dietary resources. (Food and Agricultural Organization, Rome).
- Tahira, R. and M.A. Butt. 2007. Characterization of rice bran oil. *J. Agric. Res.* 45(3):225-230.
- Tang, B., L. Feng, Y. Liu, J. Jiang, K. Hu, W. Jiang, S. Li and X. Zhou. 2012. Methionine requirement of grass carp (*Ctenopharyngodon idella*) during middle growth stage. *Chin. J. Anim. Nutr.* 24(11):2263-2271.
- Viola, S., H. Angeoni and E. Lahav. 1992a. Limits of protein sparing by lysine supplementation of pond fish feeds. *Isr. J. Aquacult-Bamid.* 44(4):125-126.
- Walton, M.J., C.B. Cowey and J.W. Adron. 1984. The effect of dietary lysine levels on growth and metabolism of rainbow trout (*Salmo gairdneri*). *Br. J. Nutr.* 52:115-122.
- Wang, S., Y.H. Liu, L.X. Tian, M.Q. Xie, H.J. Yang, Y. Wang, and G.Y. Liang. 2005. Quantitative dietary lysine requirement of juvenile grass carp *Ctenopharyngodon idella*. *Aquaculture*, 249:419-429.
- Wilson, R.P. 1985. Amino acid and protein requirements of fish. In: *Nutrition and Feeding in Fish* (Cowey, C.B., A.M. Mackie and J.G. Belleds), pp. 1-15. Academic Press, London.
- Yang, H., Y. Liu, L. Tian, G. Liang and H. Lin. 2010. Effects of Supplemental Lysine and Methionine on Growth Performance and Body Composition for Grass Carp (*Ctenopharyngodon idella*). *Am. J. Agric. Biol. Sci.* 5(2):222-227.
- Webster, C.D., L.G. Tiu, A.M. Morgan and A.L. Gannam. 2000. Differences in growth in blue catfish *Ictalurus furcatus* and channel catfish *I. punctatus* fed low-protein diets with and without supplemental methionine and/or lysine. *J. World Aquac. Soc.* 31:195-205.
- Williams, K., C. Barlow and L. Rodgers. 2001. Efficacy of crystalline and protein-bound amino acids for amino acid enrichment of diets for barramundi/Asian seabass (*Lates calcarifer* Bloch). *Aquac. Res.* 32: 415-429.
- Zarate, D.D. and R.T. Lovell. 1997. Bioavailability of free vs. protein-bound lysine in practical diets for young channel catfish (*Ictalurus punctatus*). *Aquaculture*, 159(1-2):87-100.

CONTRIBUTION OF AUTHORS

S. No.	Author name	Contribution	Signature
1.	Faryad Ali	Performed the experiment and collected the data	
2.	Muhammad Afzal	Supervised the experiment	
3.	Syed Zakir Hussain Shah	Helped in design of experiment and critically checked and improved the manuscript	
4.	Mahroze Fatima	Generated the idea, designed the study and wrote the manuscript	
5.	Khadija Akram	Helped in collection and analysis of data and writing of the manuscript	
6.	Rashid Hussain	Helped in improving the study and writing the manuscript	